

Application of the generalized three-dimensional Jordan-Wigner transformation to the bilayer Heisenberg antiferromagnet

B. Bock¹ and M. Azzouz^{1,2,*}

¹*Department of Physics, University of Northern British Columbia, Prince George, British Columbia, Canada V2N 3Z9.*

²*Max-Planck Institut für Physik Komplexer Systeme, Nöthnitzer Str.38, 01187 Dresden, Germany*
(February 6, 2008)

We extend the definition of the Jordan-Wigner transformation to three dimensions using the generalization of ideas that were used in the two-dimensional case by one of the present authors. Under this transformation, the three-dimensional XY Hamiltonian is transformed into a system of spinless fermions coupled to a gauge field with only two nonzero components. We calculate the flux per plaquette for the 3 elementary perpendicular plaquettes of a cubic lattice, and find that it is nonzero for only two of the plaquettes. We provide a simple interpretation for the average phase-per-plaquette being π on the plaquettes where it is nonzero. Then we apply these new findings to the investigation of the Heisenberg bilayer antiferromagnet.

PACS numbers: 75.10.Jm, 75.50.Ee

In 1928, Jordan and Wigner¹ introduced a transformation to write the spin operators S^- and S^z in terms of Fermi operators in one dimension. For example, this transformation allows a mapping of the XY Hamiltonian in one dimension into a Hamiltonian of noninteracting spinless fermions. Trying to generalize this result to two dimensions proved to be difficult. But, a natural extension of the Jordan-Wigner (JW) transformation was introduced in 1993 by one of the present authors² to study the interchain-coupling effect on the one-dimensional spin-1/2 Heisenberg antiferromagnets. Several other authors have given implicit ways of defining a generalized JW transformation in two dimensions.^{3,4} The generalization of the JW transformation has been attempted even in three dimensions by Huerta and Zanelli⁵ who introduced an implicit transformation, and by Kochmański⁶ whose transformation, however, does not preserve some of the spin-commutation relations.

In this work, we follow the procedure of Ref.² to extend the definition of the JW transformation in three dimensions. This paper is organized as follows. We start by reviewing the JW transformation in one and two dimensions. Then we define its three dimensional version. For the XY model, we calculate the components of the effective gauge field that results from the transformation, and the flux per plaquette as one spinless-fermion moves around it. Finally, we use this transformation to study the effect of the interlayer coupling on the Heisenberg bilayer.

The JW transformation can be performed independent of any model Hamiltonian. In one dimension this is written as

$$S_i^- = c_i e^{i\phi_i}, \quad S_i^z = c_i^\dagger c_i - 1/2. \quad (1)$$

The phase $\phi_i = \pi \sum_{j=0}^{i-1} c_j^\dagger c_j = \pi \sum_j w(i, j) n_j$ where

$$w(i, j) = \Theta(i - j)(1 - \delta_{i,j}), \quad (2)$$

with $\Theta(i - j) = 1$ for $j \leq i$ and 0 for $j > i$ being the (discrete) Heaviside step function. Note that the phase term in this transformation is obtained by summing over

all those sites to the left of site i . Also, when we calculate spin commutation relations we find that the fermion-number operator, at the site with the lowest index and hosting one spin of the commutator, is contained only once in the resulting phase (refer to Ref.² for more details). This assures that the spin commutation relations are preserved. In two dimensions, a generalized transformation based on this idea has been introduced in Ref.^[2]. This was written as:

$$S_{i,j}^- = c_{i,j} e^{i\pi\phi_{i,j}}, \quad S_{i,j}^z = c_{i,j}^\dagger c_{i,j} - 1/2, \\ \phi_{i,j} = \sum_{\alpha=0}^{i-1} \sum_{\beta=0}^{\infty} n_{\alpha,\beta} + \sum_{\beta=0}^{j-1} n_{i,\beta}, \quad (3)$$

where now a given site is specified by two indices, i along the x-axis and j along the y-axis. In this transformation, the phase at site (i, j) is obtained by summing over all sites to the left of the vertical line passing through (i, j) as well as all sites directly below (i, j) . This is a necessary and sufficient condition that allows the generalized JW transformation to preserve all spin commutation relations. The phase term in (3) can be written as: $e^{i\pi \sum_{\mathbf{z}=0}^{\infty} w(\mathbf{x}, \mathbf{z}) n(\mathbf{z})}$ with

$$w(\mathbf{x}, \mathbf{z}) = \Theta(x_1 - z_1)(1 - \delta_{x_1, z_1}) \\ + \Theta(x_2 - z_2)\delta_{x_1, z_1}(1 - \delta_{x_2, z_2}); \quad (4)$$

$\mathbf{x} = (x_1, x_2)$ and $\mathbf{z} = (z_1, z_2)$ denoting sites on the square lattice. A necessary condition for the preservation of the spin commutation relation is that the following condition is satisfied

$$e^{i\pi w(\mathbf{x}, \mathbf{z})} = -e^{i\pi w(\mathbf{z}, \mathbf{x})}. \quad (5)$$

A relation that is indeed fulfilled by Eq. (4).

Next, consider the transformation of the XY model under (3), and calculate the flux per plaquette generated as one spinless fermion moves around an elementary plaquette. The phase change is found to be:

$$(n_{i,j} - n_{i+1,j})\pi. \quad (6)$$

This result is more general than the one of Fradkin,³ who reported that the phase should be π , and also shows that Shaofeng's⁷ claim that the flux per plaquette is zero is not true.

Using the same procedure described above for one and two dimensions, we define the JW transformation in three dimensions by:

$$\begin{aligned} S_{i,j,k}^- &= c_{i,j,k} e^{i\pi\phi_{i,j,k}}, \quad S_{i,j,k}^z = c_{i,j,k}^\dagger c_{i,j,k} - 1/2 \\ \phi_{i,j,k} &= \sum_{\epsilon=0}^{k-1} \sum_{\alpha=0}^{\infty} \sum_{\beta=0}^{\infty} n_{\alpha,\beta,\epsilon} + \sum_{\alpha=0}^{i-1} \sum_{\beta=0}^{\infty} n_{\alpha,\beta,k} \\ &\quad + \sum_{\beta=0}^{j-1} n_{i,\beta,k} \end{aligned} \quad (7)$$

where now the phase is:

$$\phi(\mathbf{x}) = \sum_{z=0}^{\infty} w(\mathbf{x}, \mathbf{z}) n(\mathbf{z}) \quad (8)$$

with

$$\begin{aligned} w(\mathbf{x}, \mathbf{z}) &= \Theta(x_3 - z_3)(1 - \delta_{x_3, z_3}) \\ &\quad + \Theta(x_1 - z_1)\delta_{x_3, z_3}(1 - \delta_{x_1, z_1}) \\ &\quad + \Theta(x_2 - z_2)\delta_{x_1, z_1}\delta_{x_3, z_3}(1 - \delta_{x_2, z_2}), \end{aligned} \quad (9)$$

$\mathbf{x} = (x_1, x_2, x_3)$ and $\mathbf{z} = (z_1, z_2, z_3)$ are now sites on the cubic lattice. It is straightforward to check again that Eq. (5) is satisfied, and that all spin commutation relations are preserved. To obtain the phase (8) we choose a plane going through the site where we want to define the transformation. Here the plane chosen is the one perpendicular to the vector $(0, 0, 1)$. Then we sum over those sites on this plane as in the two-dimensional transformation. Finally we sum over all sites that are behind this plane.

As for the phase change that occurs when a spinless fermion completes a motion around a plaquette, we consider the elementary plaquettes in the xy , xz , and yz planes. The flux per plaquette in the xy plane turns out to be the same as in two dimensions, that is

$$(n_{i,j,k} - n_{i+1,j,k})\pi. \quad (10)$$

For a spinless fermion starting at site (i, j, k) , $n_{i,j,k} = 1$. Then for this fermion to go to site $(i+1, j, k)$, this later site has to be empty due to Pauli exclusion principle; thus $n_{i+1,j,k} = 0$. In this approximation one finds that the phase is π . For a plaquette in the yz plane the result is

$$(n_{i,j,k+1} - n_{i,j,k})\pi. \quad (11)$$

Similarly, the phase per plaquette can be set to be approximately π . As for the phase change around an elementary plaquette in the xz plane, it is found to be identically zero.

The results for these fluxes per plaquette suggest that as a spinless fermion moves around a given plaquette, it

will couple to a gauge field if the plaquette belongs to plane xy or yz , but no coupling to a gauge field occurs in the remaining plane xz . This thus suggests that the effective magnetic field to which the spinless fermions couple has only two components. To show this, we consider the XY model written for coupled layers:

$$\begin{aligned} H_{XY} &= \frac{J}{2} \sum_{i,j,k} S_{i,j,k}^- S_{i+1,j,k}^+ + \frac{J}{2} \sum_{i,j,k} S_{i,j,k}^- S_{i,j,k+1}^+ \\ &\quad + \frac{J_{\perp}}{2} \sum_{i,j,k} S_{i,j,k}^- S_{i,j,k}^+ + (\text{H.c.}) \end{aligned} \quad (12)$$

where J and $J_{\perp}$ are respectively the intralayer and inter-layer coupling constants. Under the JW transformation (7), the Hamiltonian (12) gives

$$\begin{aligned} H_{XY} &= J \sum_{i,j,k} c_{i,j,k}^\dagger c_{i+1,j,k} e^{i\pi\nabla_i\phi_{i,j,k}} \\ &\quad + J \sum_{i,j,k} c_{i,j,k}^\dagger c_{i,j,k+1} e^{i\pi\nabla_k\phi_{i,j,k}} \\ &\quad + J_{\perp} \sum_{i,j,k} c_{i,j,k}^\dagger c_{i,j,k} e^{i\pi\nabla_i\phi_{i,j,k}} + (\text{H.c.}) \end{aligned} \quad (13)$$

Here $\nabla_i\phi_{i,j,k} = \phi_{i+1,j,k} - \phi_{i,j,k}$ and $\nabla_k\phi_{i,j,k} = \phi_{i,j,k+1} - \phi_{i,j,k}$ designate the discrete derivatives along the x and z -axes, respectively. This effective Hamiltonian describes the motion of spinless fermions coupled to a gauge field. The later field is given by $A_i = \nabla_i\phi_{i,j,k}$ along the x -axis, $A_j = \nabla_j\phi_{i,j,k}$ along the y -axis, and $A_k = \nabla_k\phi_{i,j,k}$ along the z -axis. This leads to

$$\begin{aligned} A_i &= \pi \left[\sum_{\beta=0}^{\infty} n_{i,\beta,k} + \sum_{\beta=0}^{j-1} (n_{i+1,\beta,k} - n_{i,\beta,k}) \right], \\ A_j &= 0, \\ A_k &= \pi \left[\sum_{\alpha=0}^{i-1} \sum_{\beta=0}^{\infty} (n_{\alpha,\beta,k+1} - n_{\alpha,\beta,k}) \right. \\ &\quad \left. + \sum_{\beta=0}^{j-1} (n_{i,\beta,k+1} - n_{i,\beta,k}) + \sum_{\alpha=0}^{\infty} \sum_{\beta=0}^{\infty} n_{\alpha,\beta,k} \right]. \end{aligned} \quad (14)$$

Note that the y -component of $\mathbf{A} = (A_i, A_j, A_k)$ is found to be identically zero. The effective magnetic field is obtained through the discrete curl of $\mathbf{A}$, that is $\mathbf{B}(i, j, k) = \nabla \times \mathbf{A}(i, j, k)$, [vectors $\mathbf{A}$ and $\mathbf{B}$ depend on the position (i, j, k)]. The components of $\mathbf{B}$ which are given by

$$\begin{aligned} B_i &= \pi(n_{i,j,k+1} - n_{i,j,k}), \\ B_j &= 0, \\ B_k &= \pi(n_{i,j,k} - n_{i+1,j,k}), \end{aligned} \quad (15)$$

are consistent with the results of the fluxes per plaquette in Eqs. (10) and (11), (the unit area being 1). Therefore the effective magnetic field to which the spinless fermions are coupled has two nonzero components only.

In the second part of this paper we apply the above findings to the study of the Heisenberg bilayer antiferromagnet that has received a lot of attention in the past

few years.⁸⁻¹⁸ The use of the JW transformation and the bond-mean-field theory^{2,19} is advantageous in comparison with the Schwinger boson approach,^{8,13,14} modified spin-wave approaches,¹¹ or the bond-operator mean-field theory,^{17,18} because no constraint on the number of particle per site is required for the JW spinless fermions. We would like to draw the attention to the bond-mean field approach^{2,19} which has so far given very good results in the case of the Heisenberg ladder,²⁰ and coupled Heisenberg ladders.²¹ The bilayer antiferromagnet is a system that presents an order-disorder second order quantum transition at zero temperature, and may be of some relevance to some of the high temperature superconductors in their low-doping normal state.²²

The Heisenberg model on a bilayer reads as follows

$$H = H_{XY} + J \sum_{i,j,k} S_{i,j,k}^z S_{i+1,j,k}^z + J \sum_{i,j,k} S_{i,j,k}^z S_{i,j+1,k}^z + J_{\perp} \sum_{i,j,k} S_{i,j,k}^z S_{i,j,k+1}^z. \quad (16)$$

Hamiltonian (16) is simplified using the approximation where the average flux per elementary plaquette is π on the xy and yz planes, and zero on the xz plane (as discussed above). To achieve this, we choose the following configuration: the phases are alternated $\dots\pi-0-\pi-0\dots$ along the adjacent bonds on the y -axis, and zero on all bonds on the remaining axes. Note that the alternated phases could be put rather on the x -axis without changing the physical results because of gauge invariance. In the case of the bilayer system, the alternated phases cannot however be put on the z -axis because there is only one bond along that axis. Further simplification is done by using the bipartite character due to antiferromagnetic correlations. Then we decouple the interacting quartic terms by introducing the alternated magnetization parameter $m_i = 2\langle S_{i,j,k}^z \rangle = 2\langle c_{i,j,k}^{\dagger} c_{i,j,k} \rangle - 1 = (-1)^i m$, and the bond parameters $Q = \langle c_{i,j,k} c_{i+1,j,k}^{\dagger} \rangle = \langle c_{i,j,k} c_{i,j+1,k}^{\dagger} \rangle$ within the planes and $P = \langle c_{i,j,k} c_{i,j,k+1}^{\dagger} \rangle$ perpendicular to the planes. The Hamiltonian takes the following form in the reciprocal space:

$$H = \sum_{\mathbf{k}} (2J + J_{\perp}) m (e_{\mathbf{k}}^{\dagger} e_{\mathbf{k}} - f_{\mathbf{k}}^{\dagger} f_{\mathbf{k}}) + \sum_{\mathbf{k}} [iJ_1 \sin k_x + \gamma(k_y, k_z)] e_{\mathbf{k}}^{\dagger} f_{\mathbf{k}} + \text{H.c.} \quad (17)$$

where $\gamma(k_y, k_z) = J_1 \cos k_y + J_{\perp 1} \cos k_z$, $e_{\mathbf{k}} = c_{\mathbf{k}}^A$ (resp. $f_{\mathbf{k}} = c_{\mathbf{k}}^B$) is the fermion operator on the sublattice A (resp. B). The quantities J_1 and $J_{\perp 1}$ are given by $J_1 = J(1 + 2Q)$ and $J_{\perp 1} = J_{\perp}(1 + 2P)$. The energy spectrum takes the expression:

$$E_{\pm}(k) = \pm \{ (2J + J_{\perp})^2 m^2 + J_1^2 \sin^2 k_x + \gamma^2(k_y, k_z) \}^{1/2} \quad (18)$$

leading to the ground-state energy per site:

$$E_{GS} = \frac{2J + J_{\perp}}{4} m^2 + 2JQ^2 + J_{\perp}P^2 - \int \frac{(d^3k)}{2} E_{+} \quad (19)$$

where by definition $(d^3k) \equiv d\mathbf{k}/(2\pi)^3$. Because of the periodic boundary conditions along the z -axis, $J_{\perp}$ is counted twice. Thus we have to divide the value of $J_{\perp}$ by 2 in all our equations, or multiply $J_{\perp}$ by two in the final results wherever $J_{\perp}$ appears.




FIG. 1. The parameters m , Q and P are plotted as a function of $J_{\perp}$. Full lines are obtained for finite m , and dashed lines are for $m = 0$. The dashed-dotted line is E_g defined by Eq. (21).

The parameters m , Q , and P are calculated self-consistently through the equations:

$$\begin{aligned} m &= \int (d^3k) m (2J + J_{\perp}) / E_{+} \\ Q &= \int (d^3k) \{ J_1 \sin^2 k_x + \gamma(k_y, k_z) \cos k_y \} / 4E_{+} \\ P &= \int (d^3k) \gamma(k_y, k_z) \cos k_z / 2E_{+}, \end{aligned} \quad (20)$$

obtained by minimizing the ground-state energy with respect to m , Q and P . Two solutions for the magnetization $m/2$ are possible depending on the value of $J_{\perp}$, Fig. 1. Let us first analyze the trivial solution $m = 0$ of Eqs. (20). In this case the energy spectrum $E_k = \pm \{ J_1^2 \sin^2 k_x + \gamma^2(k_y, k_z) \}^{1/2}$ presents an energy gap at $\mathbf{k} = (k_0, \pi, 0)$ or $(k_0, 0, \pi)$, ($k_0 = 0, \pi$):

$$E_g = |J_{\perp 1} - J_1| = |J_{\perp}(1 + 2P) - J(1 + 2Q)|, \quad (21)$$

if $J_{\perp 1} > J_1$. This condition is deduced from $\gamma(k_y, k_z) = 0$ which is essential for a vanishing gap. Numerically we find that the value above which an energy gap opens is $J_{\perp} = 2 \times 0.78J = 1.56J$, see Fig. 1. However, this does not mean that this is the critical value for the order-disorder second-order quantum transition. As for the finite m solution, figure 1 shows that $m \neq 0$ for $J_{\perp} < J_{\perp c} = 2 \times 2.1J = 4.2J$. The phase is ordered antiferromagnetically for $J_{\perp} < J_{\perp c}$, whereas it is a disordered quantum state for $J_{\perp} > J_{\perp c}$. The value $J_{\perp c} = 4.2J$ compares well to the result $(4.48J)$ of the Schwinger boson approach^{8,13,14} and that (4.3) of the self-consistent spin-wave theory.¹⁵ But it disagrees with the Monte Carlo simulations¹⁰ result $(2.5J)$ or the series

expansion²² calculation ($2.56J$). Chubokov and Morr¹¹ reported that while transverse spin-wave excitations are gapless (Goldstone modes) longitudinal spin-wave excitations are gapped for $J_{\perp c} > J_{\perp} > 0$. Our result seems to indicate that a finite value of the interlayer coupling ($J_{\perp} = 1.56J$) would be needed for longitudinal fluctuations to become relevant. Note however that we have to consider gaussian fluctuations about our mean-field point to account for the spin-wave excitations in the ordered phase. We expect that fluctuations beyond the mean-field point will lead to a better estimate of $J_{\perp c}$. The magnetization increases with $J_{\perp}$, passes through a maximum, then decreases in agreement with previous results.¹¹

As for the parameters P and Q , when $J_{\perp}$ increases P saturates at 0.5, whereas Q decreases to 0, Fig. 1. For very large values of $J_{\perp}$ this approach becomes less accurate. Perturbation expansion about the transverse dimers may be used.^{16,22}




FIG. 2. The reduced ground-state energy is plotted as a function of the reduced interlayer coupling as defined in the text. The dashed line is from Ref. [16].

Finally, the result of the calculation of the ground-state energy, E_{GS} , is reported in Fig. 2 in order to support the validity of the present approach. To compare with the result of Weihong¹⁶ obtained using Ising and dimer expansions, $\nu = E_{GS}/(4J + J_{\perp})$ is drawn as a function of $\eta = J_{\perp}/(J + J_{\perp})$. The maximum of ν at $\eta = 0.5$ coincides with that reported by Weihong, and in general both results agree very well.

In summary, we extended the definition of the JW transformation to three dimensions and gave a full analysis of its consequences in the case of the XY model. This model is found to be transformed to a system of spinless fermions that are coupled to a gauge field with two components only (one of the three components being zero). Accordingly, the flux per plaquette is nonzero for 2 elementary perpendicular plaquettes, but is zero for the third one. We gave a simple interpretation of the average flux being π for those plaquettes where it is nonzero. These findings were used to study the Heisenberg bilayer antiferromagnet within the bond-mean field theory. We recalculated the order-disorder critical coupling and the ground-state energy. Very good agreement with previous results is found. Finally, the present three-

dimensional JW transformation can be applied to models of three-dimensional quantum spin systems. For simplicity a cubic Bravais lattice was used in this work, but the transformation introduced here can be easily generalized to other Bravais lattices.

B.B. acknowledges receipt of an undergraduate NSERC award. M.A. is grateful to Prof. Shегelski for his support and the useful discussions they had together, and to Prof. Hussein for his support. He thanks Prof. Thalmeier and Dr. Yuan for the helpful discussions they had during his stay at the Max-Planck Institute in Dresden. He is also grateful to this institute for the financial support.

* Correspondance could be sent to azzouz@unbc.ca

- ¹ P. Jordan and E. Wigner, Z. Physik **47**, 631 (1928)
- ² M. Azzouz, Phys. Rev. B **48**, 6136 (1993)
- ³ E. Fradkin, Phys. Rev. Lett. **63**, 322 (1989)
- ⁴ Y. R. Wang, Phys. Rev. B **46**, 151 (1992), Phys. Rev. B **45**, 12604 (1992), and Phys. Rev. B **43**, 3786 (1991)
- ⁵ L. Huerta and J. Zanelli, Phys. Rev. Lett. **71**, 362 (1993)
- ⁶ M. S. Kochmański, J. Tech. Phys. **36**, 485 (1995)
- ⁷ W. Shaofeng, Phys. Rev. E **51**, 1004 (1995)
- ⁸ A. J. Millis and H. Monien, Phys. Rev. Lett. **70**, 2810 (1993) and Phys. Rev. B **50**, 16606 (1994)
- ⁹ A. W. Sandvik and D. J. Scalapino, Phys. Rev. Lett. **72**, 2777 (1994)
- ¹⁰ A. W. Sandvik, A. V. Chubukov, and S. Sachdev, Phys. Rev. B **51**, 16483 (1995)
- ¹¹ A. V. Chubukov and D. K. Morr, Phys. Rev. B **52**, 3521 (1995)
- ¹² A. W. Sandvik and D. J. Scalapino Phys. Rev. B **R526** 1996
- ¹³ T. Miyazaki, I. Nakamura, and D. Yoshioka, Phys. Rev. B **53**, 12206 (1996)
- ¹⁴ K.-K. Ng, F. C. Zhang, and M. Ma, Phys. Rev. B **53**, 12196 (1996)
- ¹⁵ K. Hida J. Phys. Soc. Jpn. **59**, 2230 (1990). Note that a modified spin wave theory (Ref.¹¹) gives $J_{\perp c} = 2.73J$.
- ¹⁶ Z. Weihong, Phys. Rev. B **55**, 12267 (1997)
- ¹⁷ Y. Matsushita, M. P. Gelfand, and C. Ishii, J. Phys. Soc. Jpn. **68**, 247 (1999)
- ¹⁸ D.-K. Yu, Q. Gu, H.-T. Wang, and J.-L. Shen, Phys. Rev. B **59**, 111 (1999)
- ¹⁹ M. Azzouz and C. Bourbonnais, Phys. Rev. B **53**, 5090 (1996)
- ²⁰ M. Azzouz, L. Chen, and S. Moukouri, Phys. Rev. B **50**, 6233 (1994)
- ²¹ M. Azzouz, B. Dumoulin, A. Benyoussef, Phys. Rev. B **55**, R11957 (1997)
- ²² K. Hida, J. Phys. Soc. Jpn. **61**, 1013 (1992)